



24(c) Special Local Need Labeling

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for Additional Precautionary Statements

Keep out of the reach of children, pets, wildlife, and domestic animals.

Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Wear rubber gloves when handling this product. If swallowed by human beings, domestic animals or pets this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K₁, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION
IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability. Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Important: Before using this product read and carefully observe all applicable directions, restrictions and precautionary statements on the EPA-Registered container label.
WEAR RUBBER GLOVES WHEN HANDLING THIS PRODUCT.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

24© Registrant:
Manufactured By:
HACCO, Inc.
P.O. Box 7190, Madison, WI 53707

FOR DISTRIBUTION AND USE ONLY
WITHIN THE STATE OF MICHIGAN

EPA Reg. No. 61282-46
EPA SLN No. MI-840012

Expires: Dec. 31, 2005

Application Directions Dormant bearing and non-bearing Tree Fruit Orchards and Christmas Tree Plantations.

ORCHARD MICE	RATE/Acre of RAMIK	LIMITATIONS/ DIRECTIONS
MEADOW VOLE	AERIAL OR GROUND BROADCAST TREATMENT Apply 20 lbs. of pellets per acre in two (2) appli- cations of 10 lbs. each. Applications can be made 20-40 days apart.	DORMANT APPLICATION ONLY DO NOT PILE THE BAIT Do not graze meat or dairy cattle in treated fields. Do not cut hay from treated areas and feed or use for bedding. Application is not to be made until all crops are harvested and all harvestable drops are picked up.
PINE VOLE	HAND BAIT TREATMENT Apply 20 lbs. of pellets per acre in two applications of 10 lbs. each. Applications can be made 20-40 days apart. For each application make at least two (2) bait placements per tree, by dispersing ¼ lb. of Ramik pellets per tree to active vole trails and holes (40 tree-acre planting). WHERE A MIXED POPULATION IS PRESENT, APPLY AS DIRECTED FOR PINE VOLE.	DORMANT APPLICATION ONLY Restore treated area with thatch or other vegetative debris. DO NOT PILE THE BAIT Do not graze meat or dairy cattle in treated fields. Do not cut hay from treated areas and feed or use for bedding. Application is not to be made until all crops are harvested and all harvestable drops are picked up.

**Read the Notice on the Container Before Buying or Using. If
terms are not acceptable, return at once unopened.**

Code RN.NU